



A new method for measuring N_2 emissions from denitrification in soils

Wendy Yang

Whendee Silver Lab

*Dept. of Environmental Science, Policy, and Management
University of California, Berkeley*

DOE GCEP End-of-Summer Workshop
August 14, 2007

N_2O is produced from nitrous oxide, a greenhouse gas and catalyst for stratospheric ozone depletion.





Figure from Stein and Yung 2003

Human activity accounts for over half of global nitrous oxide emissions.



Adapted from Stein and Yung 2003

Nitrogen deposition rates are projected to increase dramatically in the tropics.



Existing methods for measuring N₂ emissions from soils are flawed.

- (1) ¹⁵N tracer
 - high detection limit
- (2) Acetylene inhibition
 - underestimates N₂ production rates
- (3) Incubation in N₂ free headspace
 - difficult to keep out atmospheric N₂



N_2/Ar requires high precision to detect N_2 fluxes from upland soils.


$$\delta N_2/Ar =$$

$$\left[\left(N_2/Ar_{\text{sample}} \right) / \left(N_2/Ar_{\text{ref}} \right) - 1 \right] * 10^6$$

20 per meg change in $\delta N_2/Ar =$
10-20 ng-N/cm²/h or 2-4 ng-N/gdw/h



Liquid helium is used to quantitatively transfer gas samples.




Gas handling causes most of the variability in repeated measurements.



Biological N₂ flux must be separated from physical effects on N₂/Ar.

- Water vapor flux fractionation
- Thermal diffusion




**$\delta^{40}\text{Ar}$ could constrain mass
dependent fractionation effects on
 $\delta\text{N}_2/\text{Ar}$.**



Thermal diffusion may not be a large factor influencing $\delta N_2/Ar$.

Temperature increase	% water-filled pore space	<u>Change in $\delta N_2/Ar$ (per meg)</u>	
		0-20 cm	0-50 cm
0.5°C	30	38	61
	50	75	131
	70	123	242
0.1°C	30	8	12
	50	15	27
	70	25	49

**Conclusion:
Perseverance will be rewarded.**



Acknowledgements

Collaborators

Mark Conrad (LBNL)

Mary Firestone (UCB, LBNL)

Ralph Keeling (UC-San Diego)

Jeff Severinghaus (UC-San Diego)


Funding

DOE GCEP GREF

NSF Graduate Research Fellowship

Berkeley Atmospheric Sciences Center
Graduate Fellowship

Low nitrate availability drives nitrous oxide reduction to N₂.



aerobic microbial process

anaerobic microbial process

N₂ could be an important fate of anthropogenic nitrogen inputs.

